

石墨烯的一些多项式和基于度的拓扑指数

贾米尔·艾哈迈德¹, 穆罕默德·拉希德¹, 瓦西姆·哈立德², 阿德南·阿斯拉姆^{3*}

(1. 明哈吉大学 数学系, 拉合尔 54000;

2. 拉合尔大学 数学系, 巴格伯登 54000;

3. 工程科技大学 自然科学与人文系, 拉合尔 54000)

摘要: 图 G 的拓扑指数是与 G 相关的数值参数, 其刻画了图的拓扑并在图同构下保持不变. 而化学化合物的性质和拓扑指数是相关的. 因此, 计算了石墨烯的第 1 类和第 2 类萨格勒布多项式, 以及遗忘多项式, 同时还得到石墨烯的第 1 类和第 2 类乘法萨格勒布指数和遗忘指数.

关键词: 理论化学; 石墨烯; 萨格勒布多项式; 萨格勒布指数

中图分类号: O157.6 **文献标识码:** A **文章编号:** 1674 - 5639 (2018) 06 - 0048 - 06

DOI: 10.14091/j.cnki.kmxyxb.2018.06.010

Some Polynomials and Degree-based Topological Indices of Graphene

JAMIL Ahmad¹, MUHAMMAD Rashid¹, WASEEM Khaild², ADNAN Aslam^{3*}

(1. Department of Mathematics, Minhaj University, Lahore, Pakistan 54000;

2. Department of Mathematics, University of Lahore, Pakpattan, Pakistan 54000;

3. Department of Natural Sciences and Humanities, University of Engineering and Technology, Lahore, Pakistan 54000)

Abstract: The topological index of graph G is a numerical parameter related to G , which characterizes its topology and is preserved under isomorphism of graphs. Properties of the chemical compounds and topological indices are correlated. So we compute the first and second Zagreb polynomials and forgotten polynomial of graphene. At the same time, we get the first and second multiple Zagreb indices and forgotten index of grapheme.

Key words: theoretical chemistry; grapheme; Zagreb polynomial; Zagreb index

1 Basic Knowledge

Chemical reaction network theory is an area of applied mathematics that attempts to model the behavior of real world chemical systems. Since its foundation in the 1960s, it has attracted a growing research community, mainly due to its applications in biochemistry and theoretical chemistry. It has also attracted interest from pure mathematicians due to the interesting problems that arise from the mathematical patterns in structures of material.

Cheminformatics is an emerging field in which quantitative structure-activity (QSAR) and Structure-property (QSPR) relationships predict the biological activities and properties of nanomaterial. In these studies, some physico-chemical properties and topological indices are used to predict bioactivity of the chemical compounds^[1-5].

The branch of chemistry which deals with the chemical structures with the help of mathematical tools is called mathematical chemistry. Chemical graph theory is the branch of mathematical chemistry that applies graph theory to

收稿日期: 2018 - 09 - 25

作者简介: 贾米尔·艾哈迈德, 男, 巴基斯坦人, 助理教授, 博士, 主要从事图论和理论化学研究.

* 通讯作者: 阿德南·阿斯拉姆, 男, 巴基斯坦人, 副教授, 博士, 主要从事图论和交换代数研究, E-mail: adnan-aslam15@yahoo.com.

mathematical modeling of chemical phenomena. In chemical graph theory, a molecular graph is a simple graph (having no loops and multiple edges) in which atoms and chemical bonds between them are represented by vertices and edges respectively. A graph G with vertex set $V(G)$ and edge set $E(G)$ is connected if there exists a connection between any pair of vertices in G . The distance between two vertices u and v is denoted as $d(u, v)$ and is the length of shortest path between u and v in graph G . The number of vertices of G , adjacent to a given vertex v , is the “degree” of this vertex and will be denoted by dv . For details on basics of graph theory, any standard text such as^[6] can be of great help.

The first and the second Zagreb indices (cf. ^[7]) are defined as

$$M_1(G, x) = \sum_{uv \in E(G)} x^{[d(u) + d(v)]},$$

$$M_2(G, x) = \sum_{uv \in E(G)} x^{[d(u)d(v)]}.$$

The multiplicative version of the first and second Zagreb indices were first defined by Ghorbani and Azimi^[8] as

$$PM_1(G) = \prod_{uv \in E(G)} [d(u) + d(v)],$$

$$PM_2(G) = \prod_{uv \in E(G)} [d(u)d(v)].$$

And Recently in 2015 Furtula and Gutman^[9] introduced another topological index called forgotten index or F -index

$$F(G) = \sum_{uv \in E(G)} [d(u)^2 + d(v)^2].$$

The forgotten polynomial is defined as

$$F(G, x) = \sum_{uv \in E(G)} x^{[d(u)^2 + d(v)^2]}.$$

Graphene is an atomic scale honeycomb lattice made of carbon atoms. Graphene is 200 times stronger than steel, one million times thinner than a human hair, and world's most conductive material. So it has captured the attention of scientists, researchers, and industrialists worldwide. It is one of the most promising nanomaterials because of its unique combination of superb properties, which opens a way for its exploitation in a wide spectrum of applications ranging from electronics to optics, sensors, and biodevices. Also it is the most effective material for electromagnetic interference (EMI) shielding. Now we focus on computation of topological indices of Graphene. In Figure 1, the molecular graph of Graphene is shown.




Figure 1 Molecular graph of graphene

2 Main Results and Proofs

In this paper we compute Zagreb polynomials, forgotten polynomial, multiple Zagreb indices and forgotten index of graphene. For details about topological indices and applications we refer^[10-12] and references therein.

Theorem 1 Let G be molecular graph of graphene having t rows of Benzene rings with s Benzene ring in each row. Then the first Zagreb polynomial is

$$M_1(G, x) = (t+4)x^4 + (4s+2t-4)x^5 + (3ts-2s-t-1)x^6.$$

Proof Let G be molecular graph of grapheme (see Figure 1), then the edges of graphene are divided into three types:

$$\begin{aligned} E_{|2,2|}(G) &= \{uv \in E(G) : d(u) = d(v) = 2\}; \\ E_{|2,3|}(G) &= \{uv \in E(G) : d(u) = 2, d(v) = 3\}; \\ E_{|3,3|}(G) &= \{uv \in E(G) : d(u) = d(v) = 3\}. \end{aligned}$$

Such that

$$|E_{|2,2|}| = t+4, |E_{|2,3|}| = 4s+2t-4,$$

and

$$|E_{|3,3|}| = 3ts-2s-t-1.$$

Now from the definition, the first Zagreb polynomial of G is

$$\begin{aligned} M_1(G, x) &= \sum_{uv \in E(G)} x^{[d(u)+d(v)]} = |E_{|2,2|}|x^{2+2} + |E_{|2,3|}|x^{2+3} + |E_{|3,3|}|x^{3+3} \\ &= (t+4)x^4 + (4s+2t-4)x^5 + (3ts-2s-t-1)x^6. \end{aligned}$$

Some plots of the first Zagreb polynomial of graph structure described in Theorem 1 can refer to Figure 2.




Figure 2 Plot of first Zagreb polynomial

Theorem 2 Let G be molecular graph of graphene with t rows of Benzene rings with s Benzene ring in each row. Then the second Zagreb index is.

$$M_2(G, x) = (t+4)x^4 + (4s+2t-4)x^6 + (3ts-2s-t-1)x^9.$$

Proof From the definition of second Zagreb polynomial, we have

$$\begin{aligned} M_2(G, x) &= \sum_{uv \in E(G)} x^{[d(u) \times d(v)]} = |E_{|2,2|}|x^{2 \times 2} + |E_{|2,3|}|x^{2 \times 3} + |E_{|3,3|}|x^{3 \times 3} \\ &= (t+4)x^4 + (4s+2t-4)x^6 + (3ts-2s-t-1)x^9. \end{aligned}$$

Figure 3 below shows some plots of the second Zagreb polynomial of graph structure described in Theorem 2.

Theorem 3 Let G be molecular graph of graphene having t rows of Benzene rings with s Benzene ring in each row. Then the forgotten polynomial is

$$F(G, x) = (t+4)x^8 + (4s+2t-4)x^{13} + (3ts-2s-t-1)x^{18}.$$

Proof Form the definition of forgotten polynomial, we infer

$$\begin{aligned} F(G, x) &= \sum_{uv \in E(G)} x^{[d(u)^2 + d(v)^2]} = |E_{|2,2|}|x^{[(2)^2 + (2)^2]} + |E_{|2,3|}|x^{[(2)^2 + (3)^2]} + |E_{|3,3|}|x^{[(3)^2 + (3)^2]} \\ &= (t+4)x^8 + (4s+2t-4)x^{13} + (3ts-2s-t-1)x^{18}. \end{aligned}$$

Figure 4 below manifests several plots of the forgotten polynomial of graph structure described in Theorem 3.




Figure 3 Plot of second Zagreb polynomial




Figure 4 Plot of forgotten polynomial

Theorem 4 Let G be molecular graph of graphene having t rows of Benzene rings with s Benzene ring in each row. Then the first multiple Zagreb index is

$$PM_1(G) = (4)^{t+4} \times (5)^{4s+2t-4} \times (6)^{3ts-2s-t-1}.$$

Proof From the definition of first multiple Zagreb index

$$\begin{aligned} PM_1(G) &= \prod_{uv \in E(G)} [d(u) + d(v)] \\ &= [d(u) + d(v)]^{|E_{[2,2]}|} \times [d(u) + d(v)]^{|E_{[2,3]}|} \times [d(u) + d(v)]^{|E_{[3,3]}|} \\ &= (2+2)^{t+4} \times (2+3)^{4s+2t-4} \times (3+3)^{3ts-2s-t-1} \\ &= (4)^{t+4} \times (5)^{4s+2t-4} \times (6)^{3ts-2s-t-1}. \end{aligned}$$

Also, Figure 5 presents some plots of the first multiple Zagreb index of graph structure described in Theorem 4.

Theorem 5 Let G be molecular graph of grapheme graphene having t rows of Benzene rings with s Benzene ring in each row. Then the second multiple Zagreb polynomial is

$$PM_2(G) = (4)^{t+4} \times (6)^{4s+2t-4} \times (9)^{3ts-2s-t-1}.$$

Proof From the definition of second multiple Zagreb index, we deduce

$$\begin{aligned} PM_2(G) &= \prod_{uv \in E(G)} [d(u) \times d(v)] \\ &= [d(u) \times d(v)]^{|E_{[2,2]}|} \times [d(u) \times d(v)]^{|E_{[2,3]}|} \times [d(u) \times d(v)]^{|E_{[3,3]}|} \\ &= (2 \times 2)^{t+4} \times (2 \times 3)^{4s+2t-4} \times (3 \times 3)^{3ts-2s-t-1} \\ &= (4)^{t+4} \times (6)^{4s+2t-4} \times (9)^{3ts-2s-t-1}. \end{aligned}$$




Figure 5 Plot of first multiplicative Zagreb index
(for $t=1$ left and for $s=1$ right)

Again, Figure 6 presents some plots of the second multiple Zagreb index of graph structure described in Theorem 5.




Figure 6 Plot of second multiplicative Zagreb index
(for $t=1$ left and for $s=1$ right)

Theorem 6 Let G be molecular graph of graphene having t rows of Benzene rings with s Benzene ring in each row. Then the forgotten index is

$$F(G) = 54ts + 16t + 16s - 38.$$

Proof From the definition of forgotten index, we get

$$\begin{aligned} F(G) &= \sum_{uv \in E(G)} [d(u)^2 + d(v)^2] \\ &= |E_{\{2,2\}}| [2^2 + 2^2] + |E_{\{2,3\}}| [2^2 + 3^2] + |E_{\{3,3\}}| [3^2 + 3^2] \\ &= 8(t+4) + 13(4s+2t-4) + 18(3ts-2s-t-1) \\ &= 54ts + 16t + 16s - 38. \end{aligned}$$

Finally, in Figure 7 we show some plots of the forgotten index of graph structure described in Theorem 6.




Figure 7 Plot of forgotten index (for $t=1$ left and for $s=1$ right)

3 Conclusion

In this report, we computed some degree-based topological indices for graphene. We also computed some algebraic polynomials that describe molecular topology of graphene. Moreover we plotted our results to see the dependence on the involved structural parameter.

[References]

- [1] RUCKER G, RUCKER C. On topological indices, boiling points, and cycloalkanes [J]. Journal of Chemical Information and Computer Sciences, 1999, 39: 788–802.
- [2] SARDAR M S, ZAFAR S, FARAHANI M R. The generalized zagreb index of capra-designed planar benzenoid series $Ca_k(C_6)$ [J]. Open Journal of Mathematical Sciences, 2017, 1 (1): 44–51.
- [3] MUTEER R H, SARDAR R, RAZA A. Computing topological indices of hex board and its line graph [J]. Open Journal of Mathematical Sciences, 2017, 1 (1): 62–71.
- [4] DE N. Hyper zagreb index of bridge and chain graphs [J]. Open Journal of Mathematical Sciences, 2018, 2 (1): 1–17.
- [5] GAO W, BATSHA M B, NAZEER W. K-banhatti and K-hyper banhatti indices of dominating david derived network [J]. Open Journal of Mathematical Analysis, 2017 (1): 13–24.
- [6] WEST D B. An introduction to graph theory [M]. Englewood Cliffs: Prentice-Hall, 1996.
- [7] GUTMAN I, DAS K C. The first Zagreb index 30 years after [J]. Match Communications in Mathematical and in Computer Chemistry, 2004, 50: 83–92.
- [8] GHORBANI M, AZIMI N. Note on multiple Zagreb indices [J]. Iranian Journal of Mathematical Chemistry, 2012, 3 (2): 137–143.
- [9] FURTULA B, GUTMAN I. A forgotten topological index [J]. Journal of Mathematical Chemistry, 2015, 53: 1184–1190.
- [10] RIAZ M, GAO W, BAIG A Q. M-polynomials and degree-based topological indices of some families of convex polytopes [J]. Open Journal of Mathematical Sciences, 2018, 2 (1): 18–28.
- [11] SARDAR M S, PAN X F, GAO W, et al. Computing sanskruti index of titania nanotubes [J]. Open Journal of Mathematical Sciences, 2017, 1 (1): 126–131.
- [12] TANG Z H, LIANG L, GAO W. Wiener polarity index of quasi-tree molecular structures [J]. Open Journal of Mathematical Sciences, 2018, 2 (1): 73–83.

